

Table 2.5.12. Adult/Adolescent AIDS Cases by Single and Multiple Risk Factor Categories, Diagnosed Through 1995, Tennessee

| Patient Risk Factor(s) | Cases Having only Factors Listed | |
|--|----------------------------------|--------------|
| | Number of Cases | Percentage |
| Number of factors = 1 | | |
| Men Who Have Sex With Men (MSM) | 3148 | 61.9 |
| Injecting drug user | 488 | 9.6 |
| Persons with hemophilia, adult pediatric | 91 | 1.8 |
| Heterosexual contact | 490 | 9.6 |
| Transfusion recipient | 76 | 1.5 |
| Single Risk Factor Subtotal | 4,398 | 84.4 |
| Number of factors = 2 | | |
| MSM + Injecting drug user | 380 | 7.5 |
| MSM + Persons with hemophilia | 3 | 0.1 |
| MSM + Heterosexual contact | 69 | 1.4 |
| MSM + Transfusion recipient | 27 | 0.5 |
| Injecting drug user + Persons with hemophilia | 4 | 0.1 |
| Injecting drug user + Heterosexual contact | 132 | 2.6 |
| Injecting drug user + Transfusion recipient | 8 | 0.2 |
| Persons with hemophilia + Heterosexual contact | 1 | -- |
| Persons with hemophilia + Transfusion recipient | 7 | 0.1 |
| Heterosexual contact + Transfusion recipient | 7 | 0.1 |
| Number of factors = 3 | | |
| MSM + Injecting drug user + Persons with Hemophilia | 1 | -- |
| MSM + Injecting drug user + Heterosexual contact | 38 | 0.7 |
| MSM + Injecting drug user + Transfusion recipient | 3 | 0.1 |
| MSM + Persons with Hemophilia + Transfusion recipient | 2 | -- |
| MSM + Heterosexual contact + Transfusion recipient | 2 | -- |
| Injecting drug user + Heterosexual contact + Transfusion recipient | 5 | 0.1 |
| Multiple Risk Factor Subtotal | 689 | 13.5 |
| Risk not Identified or Reported | 105 | 2.1 |
| Total | 5,087 | 100.0 |

Source: TDH HARS

Note: No adjustment is made for reporting lag.

Data are reported as of March 1996.